



More Than the Sum of the Parts: Satellite Aerosol Remote Sensing, and Its Relationship to Sub-orbital Measurements and Models

Ralph Kahn


NASA/Goddard Space Flight Center




Saharan Dust Storm
8-day trajectory
Beginning 07 June 1967
ESSA 5 Satellite



Prospero et al., Earth & Planet. Sci. Let. 1970



AVHRR
July 1989-June 1991
Aerosol Optical Depth



AVHRR 2-Channel GISS AOT





Fig. 6. (a) and (b) Monthly mean optical thickness (τ) and optical-thickness-weighted power exponent (α) for July 1986 derived with the benchmark atmosphere-ocean model and the standard ISCCP cloud-detection scheme. (c) and (d) As in panels (a) and (b) but with a modified cloud-detection scheme that retains only pixels with channel-5 temperatures warmer than the respective composite values.


TOMS Aerosol Index

April 13, 2001 [ACE-Asia]



- A ~35 year record
- Absorbing aerosols
- **Over land & water**
- Less sensitive to BL
- 10's of km resolution


OMI AOT and SSA Retrieval Approach



- **Aerosol reduces the TOA spectral contrast** of the Rayleigh-scattering gas atmosphere.
- Higher SSA (less absorbing) **increases the TOA Radiance, decreases contrast.**
- For absorbing aerosols, **elevation also matters.**
- Land **Surface** (even desert) is **usually dark** in the UV

TOMS Aerosol Index

1979-2001 Weekly, Zonal, 1°x1° Average Global Record







From: Torres et al, JAS, 2002

OMI Aerosol Products

AI

Extinction AOD



SSA

Absorption Optical Depth

March 01, 2005


From: Omar Torres

OMI Products

- NO₂
- Tropospheric O₃
- Aerosol Index
- Aerosol Extinction Opt. Depth
- Aerosol Absorption Opt. Depth

Note: AI complements other remote-sensing aerosol products:


- Can retrieve aerosol over or mixed with cloud
- Not sensitive to small, absorbing aerosol below 2 km





Phoenix Dust Storm 05 July 2011 Phoenix New Times

SeaWiFS – Sahara Dust over Canary Islands 06 March 1998




Station Fire near JPL, Pasadena CA August-September 2010




From: <http://hometown-pasadena.com>

***MODIS** – Fires in Alaska 01 July 2004 21:40 UTC*




*Mt. Etna **Plume Structure** 27-30 October 2002*



The NASA Earth Observing System's **Terra Satellite**

*First Light:
February 24, 2000*




Source: Terra Project Office / NASA Goddard Space Flight Center

Aerosol Retrievals –
Aerosol Amount
(Optical Depth)

MODIS


The MODerate-resolution Imaging Spectroradiometer

- NASA, Terra & Aqua
 - launches 1999, 2001
 - 705 km polar orbits, descending (10:30 a.m.) & ascending (1:30 p.m.)
- Sensor Characteristics
 - 36 spectral bands ranging from 0.41 to 14.385 μm
 - cross-track scan mirror with 2330 km swath width
 - Spatial resolutions:
 - 250 m (bands 1 - 2)
 - 500 m (bands 3 - 7)
 - 1000 m (bands 8 - 36)
 - 2% reflectance calibration accuracy
 - onboard solar diffuser & solar diffuser stability monitor




Improved over AVHRR:

- Calibration
- Spatial Resolution
- Spectral Range & # Bands





Mid-Visible
AOD
“Dark Target” + “Deep Blue”

MODIS
July 2010
Monthly Average



- Water & some Land
- Globe ~ **Every 2 days**
- ~ 10:30 AM & 1:30 PM




- **Fine/Coarse** Ratio
Over Water + AOD
- Sensitivity to **PM10**

MODIS Aerosol Retrieval Algorithm Steps

MODIS Aerosol Over Ocean Algorithm

All procedures applied to individual boxes of 20 x 20 pixels at 500 m resolution (10 km at nadir)



Ocean


MODIS Over Land Algorithm

All procedures applied to individual boxes of 20 x 20 pixels at 500 m resolution (10 km at nadir)

Identify and mask all water, cloudy and snow/ice pixels.

Separate all pixels that are outside of $0.01 < \rho_{2.13} \leq 0.25$



Discard brightest 50% and darkest 20% of pixels defined with $\rho_{0.66}$



Land

Remer et al., JAS 2005


Multi-angle Imaging SpectroRadiometer



<http://www-misr.jpl.nasa.gov>

- Nine CCD push-broom cameras
- Nine view angles at Earth surface:
70.5° forward to 70.5° aft
- Four spectral bands at each angle:
446, 558, 672, 866 nm
- *Studies Aerosols, Clouds, & Surface*

Eight Years of Seasonally Averaged Mid-visible Aerosol Optical Depth from MISR




...includes bright desert dust source regions

MISR Team, JPL and GSFC


MISR-AERONET AOT Comparison for 3,995 Coincidences

Stratified by expected aerosol air mass type




MISR-MODIS *Aerosol Optical Depth* Comparison

[MISR V22 vs. MODIS/Terra Collection 5; January 2006 Coincident Data]




Over-ocean regression coefficient **0.90**
Regression line slope 0.75
MODIS QC ≥ 1



Over-land regression coefficient **0.71**
Regression line slope 0.60
MODIS QC = 3

MISR-MODIS Coincident AOT ***Outlier Clusters***



Dark Blue [MISR > MODIS] – N. Africa *Mixed Dust & Smoke*


Cyan [MODIS > MISR, AOD large] – Indo-Gangetic Plain *Dark Pollution Aerosol*

Green [MODIS >> MISR] – Patagonia and N. Australia *MODIS Unscreened Bright Surface*

MISR Retrieval Status Distribution

Overall, about **15%** of Earth's surface produces successful MISR automatic aerosol retrievals

Dark blue = Ocean retrieval
Light blue = Land retrieval




Kahn, Nelson, Garay et al., TGRS, 2009

From experience with MISR & MODIS:


*For global, $\sim 1^\circ \times 1^\circ$ AOD, in general, MISR data need to be aggregated to **~ 3 -month sampling** to converge with MODIS*

MISR-MODIS-AERONET *Sampling* Differences


[Ascension Island 18 February 2005]



Sampling: **MISR**; MODIS; **AERONET**



AOT Snapshot: **ARNET** > **MISR** > **MODIS**



AERONET Time Series - Changing AOT

Kahn et al., JGR 2007

Clean, maritime aerosol air mass, but AOT changes 60% across RH boundary


Using any one of these to represent the entire region AOT --> large errors
Taken together, they give a better picture...

Sampling Effects is a continuing story...

Aerosol Retrievals – Aerosol Type

One MODIS Aerosol Type Classification:

Low AOT (blue), **High AOT+Coarse** (green), **High AOT+Fine** (red)




From: *Kaufman et al.*, JGR, 2005


Progress Toward a Global Aerosol Type Climatology

Ralph Kahn

NASA/Goddard Space Flight Center




***CALIPSO* 6-Grouping Aerosol Type Classification**




***AERONET* Aerosol Type 7-Grouping Classification**

**Four-parameter
AERONET-derived
classification:**

- $EAE_{491,863}$
- SSA_{491}
- RRI_{670}
- $dSSA_{863-491}$



**7 Groupings
 SSA_{491} vs.
Extinction ANG**




**7 Groupings
Real RI_{670} vs.
Extinction ANG**


***PARASOL* data at Forth Crete projected onto the AERONET Aerosol Type Classification**

Four-parameter AERONET-derived classification:

- $EAE_{491,863}$
- SSA_{491}
- RRI_{670}
- $dSSA_{863-491}$




**7 Groupings
 SSA_{491} vs.
Extinction ANG**




**7 Groupings
Real RI_{670} vs.
Extinction ANG**

HSRL Aerosol Type 8-Grouping Classification



**Four-parameter
AERONET-derived
classification:**

- α_{532}/β_{532}
- β_{1064}/β_{532}
- δ_{532}
- $\beta_{1064}/\delta_{532}$




Aerosol_cci

(European Consortium)


- 4 basic components
(**Dust**, **Sea Salt**, Fine-mode **Weakly** & **Strongly** Abs.)
- Reflects theoretical information content
- External mixtures with 3 mixing fractions
- Evaluation ongoing of information content
- Output (easier to validate / compare)
 - Fine mode AOD (fine mode / total mixing fraction)
 - Dust AOD (dust / total coarse mode mixing fraction)
 - [AAOD (absorption fraction in fine mode)]

cci AOD mixing (fractions) from AEROCOM Models


Fine mode fraction




Fraction of the less absorbing component in the fine mode






Fraction of dust in the coarse mode




AOD550 (not used as a priori)



Single-scattering Phase Functions for **Different Particle Properties**



Kahn et al., JGR 1998



Los Alamos Fire, New Mexico May 9, 2000



MISR 60° Forward




MISR Nadir




MISR 60° Aft

Broad Outline of MISR Aerosol/Surface Retrieval Algorithm Steps




Stage 1

- Cloud, terrain masks
- Band & Gas corrections
- Convert to Eq. Reflectance



Stage 2

- Select water or land algorithm
- Aggregate Eq. Reflectances
- Calculate EOFs if land



Stage 3

- Compare measurements with pre-calculated options
- Calculate uncertainties

Key Algorithm Inputs & Assumptions

Stage 1

- Calibration (band corrections, etc.)
- Surface Type Information (terrain type, snow mask, wind, etc.)
- Atmospheric gas concentrations (O_3 , H_2O_v , etc.)
- Ephemeris & solar input
- Cloud masking

Stage 2

- Aggregation (How to select & process the Stage 1 reflectances)
- Retrieval approach (What to assume or derive about aerosols & surface)**

Stage 3


- Aerosol components & mixtures
- Aerosol vertical distribution
- Radiative transfer code(s)
- Look-up table resolution
- Acceptance criteria
- Uncertainty metrics

** Retrieved Quantities \leftrightarrow Information Content of the data

Pre-launch Sensitivity Studies, Post-launch Validation


Smoke from Mexico -- 02 May 2002

Aerosol:
Amount
Size
Shape




Medium
Spherical
Smoke
Particles

Dust blowing off the Sahara Desert -- 6 February 2004




Large
Non-Spherical
Dust
Particles




***MISR* Aerosol Type Discrimination**




January 2007



July 2007



MISR Climatology Dust AOD



Aerosol Type Summary

- Remote-sensing can provide optical constraints interpreted as particle *Size, Shape, and Indices of Refraction*
- A further interpretative step, entailing additional assumptions, reports particle *Chemical Composition*
- Remote-sensing *sensitivity to particle properties is much more dependent than AOD on retrieval conditions*
- *Validation Data* for aerosol type are very limited
 - *Model simulations* and *In Situ measurements* can help

GRASP: Generalized Retrieval of Aerosol and Surface Properties


Open Source

No Location-Specific Assumptions on aerosol and surface

All calculation on the fly


Retrieved parameters: Surface reflectance, aerosol: AOD, SSA, aerosol height, size information, refractive index, aerosol type, etc.

Expected practical advantages: accurate even over bright surfaces, even for high AOD, and for extended set of parameters




Dubovik et al. 2011, 2014

GRASP: towards aerosol classification



Biomass Burning



Aerosol Type Validation Approach

- No “*Ground Truth*” except from Field Campaigns (*Golden Days*)
 - Unlike *Spectral AOD* (and *ANG*) from AERONET
Particle Properties derived from AERONET entail *many more assumptions*
 - *Far fewer* Satellite-AERONET Sky-scan than Direct-sun Coincidences
- *Self-consistency* Tests
 - *Qualitative*, but useful
 - *Regional* and *Temporal Behavior* (stratified) vs. *Expectation*
- *Comparisons* with AERONET proxies
 - Compare *Seasonal*, *Inter-annual* patterns *Statistically*
 - *Fine-mode Fraction* (FMF)
 - *Effective radius* (r_e) and *variance* (σ) [two modes – *issue with def. of “modes”*]
 - *Single-scattering albedo* (SSA) [for AOD₄₄₀ > 0.4; AERONET SZA > 50°]
 - *Sphericity* (“%Sph.”) [for AERONET ANG < 1.0 only – few coincidences w/AOD>0.2]



MISR Aerosol Type Discrimination

January 2007

Sahara Desert (Arid Region)



July 2007

Mean Best Estimate Optical Depth, Sahara





AOD





Mean Best Estimate Optical Depth, Sahara



Mean Best Estimate AOD Map & Histogram Distribution








#SuccMix vs. Normalized AOD & vs. Normalized Scattering Angle Range



1–10 11–20 21–30 31–40 41–50 51–62 63–70 71–74





Most Frequent Lowest Residual Aerosol Type Mixture Group, Stratified by AOD





Histograms of Lowest Residual & All Successful Aerosol Type Mixture Groups vs. AOD

Kahn & Gaitley JGR 2015

Statistical *Comparisons* with AERONET – *Solar Village*




SAMUM Campaign Morocco – June 04, 2006




MISR SAMUM Aerosol Air Masses (V19)

04 June 2006 Orbit 34369, Path 201, Blocks 65-68, 11:11 UTC




- A **dust-laden density flow in the SE** corner of the MISR swath
- **High SSA, ANG & Fraction Spherical** region SE of Quarzazate, includes Zagora

MISR SAMUM Aerosol Air Mass Validation



Falcon F-20 HSRL
- Thin layers of small, bright pollution particles




NOAA/HYSPLIT
Back Trajectories
-Source in N Algeria for sites 2, 3 (in the highlighted area) but not 1 (outside it).

Key Attributes of the MISR Version 22 Aerosol Product

- **AOT Coverage** – *Global but limited sampling* on a monthly basis
- **AOT Accuracy** – Maintained even when particle property information is poor
- **Particle Size** – *~3 groupings reliably*; quantitative results vary w/conditions
- **Particle Shape** – *spherical vs. non-spherical robust*, except for coarse dust
- **Particle SSA** – useful for *qualitative* distinctions
- **Aerosol Type Information** – diminished when $AOT < 0.15$ or 0.2
- **Particle Property Retrievals** – *improvement expected* w/algorithm upgrades
- **Aerosol Air-mass Types** – *more robust* than individual properties



PLEASE READ THE QUALITY STATEMENT!!!

... and more details are in publications referenced therein




Adapted from: Kahn, Survy. Geophys. 2012

SEAC⁴RS – MISR Overview 19 August 2013



Passive-remote-sensing **Aerosol Type** is a **Total-Column-Effective, Categorical** variable!!


Model-Based Aerosol-Type



Taylor et al., Atmosph. Environ. 2015

Where remote-sensing data are ambiguous, can *use a model to weights the options*

MISR ANG, AAOD Results *Constrained by GoCART Model*



ANG

$$\text{Diff}_{\text{ANG}} = |\alpha_{\text{MISR}} - \alpha_{\text{GOCART}}| \leq \varepsilon_{\text{ANG}}$$


$$\text{Diff}_{\text{AAOD}} = \left| \text{Fraction}_{\text{MISR_AAOD}} - \text{Fraction}_{\text{GOCART_AAOD}} \right| \leq \varepsilon_{\text{AAOD}}$$

Four years of data (2006-2009)
Seasonally averaged

Shenshen Li et al. AMT 2015

Where remote-sensing data are ambiguous, can *use a model to weights the options*

MISR ANG, AAOD Results *Constrained by GoCART Model*



AAOD

$$\text{Diff}_{\text{ANG}} = |\alpha_{\text{MISR}} - \alpha_{\text{GOCART}}| \leq \varepsilon_{\text{ANG}}$$

$$\text{Diff}_{\text{AAOD}} = \left| \text{Fraction}_{\text{MISR_AAOD}} - \text{Fraction}_{\text{GOCART_AAOD}} \right| \leq \varepsilon_{\text{AAOD}}$$


Four years of data (2006-2009)
Seasonally averaged

Shenshen Li et al. AMT 2015


Where remote-sensing data are ambiguous, can *use a model to weights the options*

A Three-way Street:


MISR & MODIS Provide Context, SEAC⁴RS Provides Detail,
and Models Complete the Picture




Story 1: Aerosol Air Mass Type Validation



Story 2: Upwind Smoke Source & Injection Height



Story 3: Regional Aerosol Characterization




- Five Aerosol Air Masses:
- Three Smoke Plumes
 - Continental Bkgnd.
 - Continental-Smoke Mix

19 August Golden Day


R. Kahn et al., in preparation

Applications –
Transports & Plume Heights


Aerosol Material Fluxes: Atlantic Dust & Asian Pollution





MODIS AOD & Type
Low AOD, Fine BioBurn, Coarse Dust



NCEP **W Wind** - MODIS **AOD**
Correlation 2.6-5 km; May-October




Dust Transport Estimate (Tg)
May-October (Top) January-April (Bot)
Kaufman et al., JGR 2005






MODIS AOD & type, **Field Campaign** aerosol properties & vertical distribution, GEOS model winds;
Compared with GOCART and GMI model Fine-particle mass fluxes *Yu et al., JGR 2008*

Constraining Aerosol Sources, Transports, & Sinks

Complementary MISR & MODIS AOD; Saharan Dust Plume over Atlantic June 19-23, 2000




Contours: AOT=0.15 (yellow); AOT=0.5 (purple)




Kalashnikova and Kahn, JGR 2008

MISR-MODIS-NAAPS (July 4, 2000)



MISR and MODIS AOD




NAAPS Dust

NAAPS dust **plume extent** predictions:


- In **qualitative agreement** with MISR & MODIS
- Magnitudes differ... constrains dust **Source Strength & Removal Rate**

CALIPSO Space Lidar

First Light - June 07, 2006




523 and 1064 nm channels; ~100m horizontal resolution



http://www.nasa.gov/mission_pages/calipso/main/index.html

Aerosol Sources, Processing, Transports, Sinks: **Lidar** + **Model**




August 2007 Saharan dust “D” and smoke “S” event
mapped by CALIPSO 532 nm backscatter, with superposed
model back trajectories and airborne HSRL observations


Piecing together the bigger picture. Consistency requires –

- An understanding of the *mechanisms* governing aerosol evolution
- Adequately constrained *initial & boundary* conditions

Changes in geometric perspective with angle






Changes in geometric perspective with angle



Saharan Dust Source Plume

Bodele Depression Chad June 3, 2005 Orbit 29038






Dust is injected near-surface...

Kahn et al., JGR 2007

Transported Dust Plume

Atlantic, off Mauritania March 4, 2004 Orbit 22399




Transported dust finds elevated layer of relative stability...


Kahn et al., JGR 2007

MISR height analysis of World Trade Center plume

12 September 2001





MISR
70°
image



MISR
stereo
heights
of plume
patches


Oregon Fire Sept 04 2003

Orbit 19753 Blks 53-55 MISR Aerosols V17, Heights V13 (no winds)




Detail of Wildfire Source Region


Oregon Fire Sept 04 2003




MISR Nadir **275 m** Image



MISR **Plume Heights** for Sub-patches





MODIS Image + **Fire Power**




Very Simple Plume Parcel Model

→ **Broad swath + high spatial resolution** needed to characterize sources

Wildfire Smoke Plume Database



<http://www-misr2.jpl.nasa.gov/EPA-Plumes/>




N. America
2002, 2004-2007

Africa
2005, 2006

D. Nelson, et al., Remt. Sens. 2013


*MISR Stereo-Derived **Plume Heights***
***07 May 2010** Orbit 55238 Path 216 Blk 40 UT 12:39*



D. Nelson and the MISR Team, JPL and GSFC

MISR Research *Aerosol Retrievals*



07 May 2010 Orbit 55238 Path 216 Blk 40 UT 12:39




- Plume Particles**
- Distinct from background
 - *larger, darker*
 - *much higher AOD*
 - *Non-spherical* dominated
 - Brighten downwind
 - Tend to decrease in size downwind

Wildfire Smoke Injection Heights & Source Strengths


[These are the two key parameters representing aerosol
sources in climate models]




Val Martin et al. ACP 2010



MODIS Smoke Plume Image & Aerosol Amount Snapshots



GoCART Model-Simulated Aerosol Amount Snapshots
for Different Assumed Source Strengths




Different Techniques for Assuming Model Source Strength
Overestimate or **Underestimate** Observation
Systematically in Different Regions

Petrenko et al., JGR 2012

Applications – Air Quality





Regional-scale **Air Quality** Assessment




Zhang et al., GRL. 2007

Surface-based mass-spec aerosol composition measurements


- Need to isolate *Near-surface Aerosol Component*
- Need sufficient *Spatial-Temporal Coverage* to capture *Severe Events*
- Detailed *Chemical Speciation* often required
- *High Spatial Resolution* often required (e.g., in Urban areas)




2002 4-day regional pollution event
Posts = EPA PM_{2.5} surface obs.
Color contours = MODIS AOT



Pollution Aerosol Concentrated in Ganges Valley near Kanpur, India (MISR)




MISR mid-visible AOD
[Winter, 2001-2004; white --> AOD >0.6]




NCEP Winds + Topography
[Black=surface; Red=850 mb;
contours=vertical, solid=subsidence]

Air Quality: BL Aerosol Concentration [MISR + MODIS] AOD & GEOS-Chem Vertical Distribution



[BL PM_{2.5}] /
[Total-col. AOD]
2001- 2006

20 40 60 80 100 120 140
 η ($\mu\text{g}/\text{m}^3$)






Derived
PM_{2.5}

0 5 10 15 20 50 80
Satellite-derived PM_{2.5} ($\mu\text{g}/\text{m}^3$)

MISR - GEOS-Chem Regression Model To Map **Near-surface Aerosol Pollution**

MISR-Constrained Model




MISR / GEOS-CHEM **Retrieval**

Surface network (IMPROVE) measurements


- Using MISR **Particle Shape** as well as AOT to constrain model --> much better result
- Can add column **Size** and **SSA** information when MISR retrieval is more robust

Characterizing seasonal changes in anthropogenic and natural aerosols w.r.t. preceding season over the Indian Subcontinent




Index uses MISR-retrieved particle shape and size constraints to separate natural from anthropogenic aerosol

Mapping AOD & Aerosol Air-Mass-Type in Urban Regions



Urban Pollution AOD & Aerosol Air Mass Type Mapping

INTEX-B, 06 & 15 March 2006




March
06

March
15

Aerosol Air Masses: *Dust* (non-spherical), *Smoke* (spherical, spectrally steep absorbing), and *Pollution* particles (spherical, spectrally flat absorbing) dominate specific regions

Applications –
Aerosol Climate Forcing

Multi-year Annual Average *Aerosol Optical Depth* from Different Measurements + *Synthesis* (S^*)



From: Kinne et al. ACP 2006

Even DARF and Anthropogenic DARF are *NOT* Solved Problems (Yet)






FIGURE SPM-2. Global-average radiative forcing (RF) estimates and ranges in 2005 for anthropogenic carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O) and other important agents and mechanisms, together with the typical geographical extent (spatial scale) of the forcing and the assessed level of scientific understanding (LOSU). The net anthropogenic radiative forcing and its range are also shown. These require summing asymmetric uncertainty estimates from the component terms, and cannot be obtained by simple addition. Additional forcing factors not included here are considered to have a very low LOSU. Volcanic aerosols contribute an additional natural forcing but are not included in this figure due to their episodic nature. Range for linear contrails does not include other possible effects of aviation on cloudiness. {2.9, Figure 2.20}

**IPCC AR3, 2001
(Pre-EOS)**

**IPCC AR4, 2007
(EOS + ~ 6 years)**


The *Current* Assessment of Climate Forcing Factors



Global average, & compared to other uncertainties in the models – **What about aerosol type?!**

The next big area to address

Measurement Synthesis: Aerosol Short-wave Direct Radiative Forcing




Outgoing zonal TOA fluxes calculated with Monte-Carlo Aerosol-Cloud-Radiation model constrained with *MISR AOD*, *AERONET* particle properties, *GOCART* interpolation, and using *four choices of cloud data* from ISCCP and CERES (hourly monthly and monthly mean).

Results are *compared to CERES* and validated using BSRN.

“Overall, such agreements suggest that global data sets of aerosols and cloud parameters released by recent satellite experiments (MISR, MODIS and CERES) *meet the required accuracy to use them as input to simulate the radiative fluxes* within instrumental errors.” -- Kim & Ramanathan JGR 2008

Aerosol Long-wave Radiative Forcing Over Desert



Terra/ **MODIS** – 1 km resolution cloud-clearing for 30 km CERES footprint

MISR – AOD over bright desert surface

CERES – TOA LW flux as a function of MISR AOD [$\sim 15 \pm 5 \text{ W}/(\text{m}^2 - \text{AOD})$]

Region 4 – Dust is near-surface [**TOMS** does not see it; LW flux indep. of AOD]

Result depends on dust vertical distribution & H₂O_v

Zhang & Christopher GRL 2003

Over-Land Aerosol Short-wave Radiative Forcing w/Consistent Data

The slope of:


TOA albedo vs. AOD

For data stratified by:

Surface BHR

Produces:


**Spectral aerosol
radiative efficiency**





Bright surface
+ dark aerosol
= decreasing
albedo w/AOD

Applications –
Long-Term AOD Trends

MODIS10-Year Global/Regional Over-Water AOD Trends



Trend



Statistical
Significance

- Statistically negligible ($\pm 0.003/\text{decade}$) *global-average* over-water AOD trend
- Statistically significant increases over the *Bay of Bengal, E. Asia coast, Arabian Sea*


Using MSIR **Products**

Introduction to Aerosol Product Content




There is a vast amount of detail -- and documentation

- I'd aim to get a good overview of the capabilities**
- Then learn what you need of the products & software**
- The Data Archives (DAACs) are set up to help users**

MODIS Granule




<http://eosweb.larc.nasa.gov/MISRBR/>



Schematic Hierarchy of MISR Aerosol-Surface Product Spatial Domains


NOT TO SCALE





Version 5 . 0

requires IDL 6 . 1





hdfscan

snapshot

GUI version 3.3 (last modified on: 21nov2001)
Core hdfscan version 3.3.1 (last modified on: 20feb2002)

Choose HDF file About hdfscan Quit

Image number: 1
Structure: Standard
Field: AveSceneElev
Number of dimensions: 3
YDim (range: 1 - 512, stride: 1)
XDim (range: 1 - 128, stride: 1)
SOMBlockDim (range: 30 - 30, stride: 1)
Number of samples per line: 512
Number of lines in image: 128

Gray scale image op

Display image no. 1 through 1 OK

Discard image no. 1 through 1 OK

Data display, output, or edit

Choose a structure name

Standard
Regional

Choose a range of fields (format)

AveSceneElev
StdDevSceneElev
StdDevSceneElevRelSlp
PtElev
GeoLatitude
GeoLongitude
SurfaceFeatureID
AveSurfNormAzAng
AveSurfNormZenAng

Table #1 : AveSceneElev

Filename: /data/bank/anc/AGP/database/path001/MISR_AM1_AGP_P001_F01_24.hdf

	Sample 1	Sample 2	Sample 3	Sample 4	Sample 5	Sample 6
Line 1	2734	2733	2732	2732	2732	2731
Line 2	2736	2735	2734	2734	2734	2733
Line 3	2738	2738	2737	2737	2736	2736
Line 4	2740	2740	2739	2739	2739	2738
Line 5	2743	2742	2742	2741	2741	2740
Line 6	2745	2745	2744	2744	2743	2743
Line 7	2748	2747	2747	2746	2746	2745
Line 8	2751	2750	2750	2750	2749	2748
Line 9	2754	2753	2753	2752	2752	2751
Line 10	2757	2756	2755	2755	2754	2754
Line 11	2760	2759	2758	2758	2757	2757
Line 12	2763	2762	2761	2761	2760	2760
Line 13	2766	2765	2764	2763	2763	2762
Line 14	2768	2768	2767	2766	2766	2765
Line 15	2771	2771	2770	2769	2768	2768
Line 16	2774	2774	2773	2772	2771	2770
Line 17	2777	2776	2776	2775	2774	2774
Line 18	2780	2780	2779	2778	2777	2776
Line 19	2783	2782	2781	2780	2779	2779
Line 20	2785	2785	2784	2783	2783	2782

Quit

Target SOM coordinates

Filename: /data/bank/anc/AGP/database/path001/MISR_AM1_AGP_P001_F01_24.hdf

*** target SOM coordinate location ***

path = 126 block = 66 latitude = 30.002 longitude = 110.001
target offset from block center = 66.6 km
1.1 km SOM grid: YDim = 317 XDim = 127
2.2 km SOM grid: YDim = 159 XDim = 64
17.6 km SOM grid: YDim = 20 XDim = 8
35.2 km SOM grid: YDim = 10 XDim = 4
70.4 km SOM grid: YDim = 5 XDim = 2

path = 125 block = 67 latitude = 29.999 longitude = 109.997
target offset from block center = -64.3 km
1.1 km SOM grid: YDim = 198 XDim = 14
2.2 km SOM grid: YDim = 99 XDim = 7
17.6 km SOM grid: YDim = 13 XDim = 1
35.2 km SOM grid: YDim = 7 XDim = 1
70.4 km SOM grid: YDim = 4 XDim = 1

closest path = 125

Print Save in file Quit

Image #1 : AveSceneElev

Filename: MISR_AM1_AGP_P001_F01_24.hdf

Min threshold 48.00
Max threshold 3036.00

Linear Equalized
Normal Inverted
Saturated Clipped

Reset display

Display histogram

Show cross-section plots

Image status: Linear, normal, saturated [48.00, 3036.00]

Line 106 Sample 179 Pixel value 2.934000e+03 Save

Choose geodetic lat-lon

Enter geodetic latitude of target point 30.0
Enter geodetic longitude of target point 110.0

OK Quit

MISR Level 2 Aerosol Product – Key Fields

RegBestEstimate**SpectralOptDepth** – 4 values, the mean of all successful models in the climatology
RegBestEstimate**SpectralOptDepthUnc** – 4 values, standard deviation of all successful models
RegBestEstimate**AngstromExponent** – slope of linear least-squares fit to log AOTs vs. log of four wavelengths
RegBestEstimate**SpectralSSA** – aggregated among successful models by binning rules
RegBestEstimateSpectral**OptDepthFraction** -- AOT fraction grouped by (1) rad < 0.35; (2) 0.35 < rad < 0.7; (3) rad > 0.7; (4) spherical; (5) non-spherical
RegBestEstimate**NumberFraction** – five fractional values, based on relative particle number concentration
RegBestEstimate**VolumeFraction** – five fractional values, volume-weighted
RegBestEstimate**QA** – 0=1 successful mixture; 1=(>1) successful mixtures; 2=not used; 3=no successful mixtures
Reg**LowestResidSpectralOptDepth**, etc. – value for mixture having lowest ChiSqAbs or ChiSqHet

AlgTypeFlag – gives the algorithm type used for the region: 1=dark water; 3=het. land, 253=fill
RegClassInd – 0=clear; 1=solar oblique; 2=topo. complex; 3=cloudy; 4=no data; 253=fill
AerRetrSuccFlag – 1=no successful mixtures; 3=algorithm failure; 5=insufficient data; 7=successful retrieval
NumSuccAerMixture – number of successful mixtures
NumCamUsed – number of cameras with good data and not in glint (max=9)
NumAcceptSubr – number of 1.1 km sub-regions within region passing all criteria for use (max=256)
RegEqRefl – observed regional equivalent reflectances used in retrieval (max=36)
SolZenAng, ScatterAng, GlitterAng, etc. – region-specific geometry
AerRetrSuccFlagPerMixture – 1=mixture successful; 251, 252, 253=mixture not successful
OptDepthPerMixture – lowest resid. AOT for each mixture, successful **or not**. (V15 max=24; V16 max=74)
ChisqAbs, Geom, Spec, Maxdev, Het – per mixture ChiSq values, whether the mixture was successful **or not**

RetrAppMask (**SubregParamsAer**) – 0=clear; 1=missing data; 2=poor quality; 7=cloudy, etc.

Reference: MISR [Data Product Specification](#) document, Section 8: Level 2 Aerosol/Surface Product

MISR Level 2 Aerosol Product – Additional Files Needed for Some Applications

- **ACP** (**Aerosol Climatology Product**)
 - **APOP** – Aerosol Physical and **Optical Properties** table – 8 Components (V16)
 - Aerosol **Mixture File** – Up to three-component mixtures (74 in V16; 24 in V15)
- **AGP** (**Ancillary Geographic Product**)
 - One for each of 233 SOM “Paths,” 1.1 km resolution
 - Pixel lat, long, surface type, elevation
- **ARP** (**Ancillary Radiometric Product**)
 - Pre-flight & in-flight radiometric calibration parameters, spectral band passes
 - Scale Factors to convert unpacked DN into Radiance (W/m²-str-micron)

Reference: MISR [Data Product Specification](#) doc, Section 9. MISR Ancillary Products

Most key aerosol quantities are **also stored in Aerosol Product metadata**

Some Satellite-Aerosol-Product-Related Web Sites

- <http://earthobservatory.nasa.gov> NASA **Earth Observatory** Imagery, Captions, Features
- <https://worldview.earthdata.nasa.gov> **Worldview Data Display** Website (MODIS, VIIRS)
- <http://daac.gsfc.nasa.gov/giovanni/> **Giovanni** on-line visualization & analysis tools
- <http://neo.sci.gsfc.nasa.gov> **NASA Earth Observations** Website
- <https://lance.modaps.eosdis.nasa.gov/realtime/2008031/> MODIS **Near-Real-Time Imagery**
- <https://earthdata.nasa.gov/earth-observation-data/near-real-time/download-nrt-data>
Near-Real-Time MODIS, MISR, AIRS, AMSR2, MLS, OMI, VIIRS data
- <http://rapidfire.sci.gsfc.nasa.gov/> **MODIS** rapid response imagery
- <http://www.goes.noaa.gov/#main-content> NOAA **GOES Geostationary Imagery**
- <http://cloudsgate2.larc.nasa.gov/index.html> NASA Langley Satellite **Imagery & Cloud** Data
- <http://cloudsgate2.larc.nasa.gov/cgi-bin/predict/predict.cgi> Satellite **Overpass Predictor**
- http://disc.sci.gsfc.nasa.gov/acdisc/TOMS/tomsl3_dataset.gd.html TOMS **O₃, Gases, Arsls**
- <http://www.ssd.noaa.gov/PS/FIRE/GASP/AOD/> NOAA **MODIS AOD** Maps
- <http://neo.sci.gsfc.nasa.gov> **CALIPSO Browse** Imagery

Some Satellite-Aerosol-Product-Related Web Sites

- <http://eosweb.larc.nasa.gov> **MISR, CERES, SAGE**, MOPITT, TES, data & docs
- <http://www-misr.jpl.nasa.gov> **MISR** Home page; background, image gallery,..
- <http://modis-atmos.gsfc.nasa.gov/IMAGES/index.html> **MODIS** global browse imagery
- <http://modis-atmos.gsfc.nasa.gov/> **MODIS** atmosphere products & docs
- <http://ncc.nesdis.noaa.gov/VIIRS/> NOAA **VIIRS** data & docs
- <http://www-modis.bu.edu/> **MODIS** land cover, surface albedo, and BRDF products
- <http://modis-fire.umd.edu/> MODIS-UMD **Fire products** & docs
- <http://activefiremaps.fs.fed.us/activefiremaps.php> MODIS global **Fire occurrence mapper**
- <http://www.star.nesdis.noaa.gov/smcd/spb/aq/> IDEA merged MODIS-EPA **Air Quality**
- <http://alg.umbc.edu/usaq/> UMBC **Air Quality** events
- <http://oceancolor.gsfc.nasa.gov/SeaWiFS/BACKGROUND/> **SeaWiFS** data & docs
- <http://aeronet.gsfc.nasa.gov/> **AERONET** AOT & properties, data & docs
- <http://mplnet.gsfc.nasa.gov> **MPLNET** Micro-pulse LIDAR data & docs

Using *In Situ* Observations
To Add Value to Satellite Data

SAM-CAAM

[Systematic Aircraft Measurements to Characterize Aerosol Air Masses]



[This is currently a *concept-development effort*, not yet a project]

Primary Objectives:

- Interpret and *enhance 16+ years of satellite aerosol retrieval* products
- *Characterize statistically particle properties* for major aerosol types globally, to provide detail unobtainable from space, but needed to *improve*:
 - Satellite aerosol *retrieval algorithms*
 - The *translation between satellite-retrieved aerosol optical properties and species-specific aerosol mass and size tracked in aerosol transport & climate models*

Kahn et al., BAMS submitted


SAM-CAAM *Concept*

[Systematic Aircraft Measurements to Characterize Aerosol Air Masses]

- *Dedicated Operational Aircraft* – routine flights, 2-3 x/week, on a continuing basis
- *Sample Aerosol Air Masses* accessible from a given base-of-operations, then move; project science team to determine schedule, possible field campaign participation
- Focus on *in situ measurements required* to characterize particle *Optical Properties*, *Chemical Type*, and *Mass Extinction Efficiency* (MEE)
- *Process Data Routinely* at central site; instrument PIs develop & deliver algorithms, upgrade as needed; data distributed via central web site
- Peer-reviewed Paper identifying *4 Payload Options*, of varying ambition; subsequent selections based on agency buy-in and available resources

SAM-CAAM is feasible because:

Unlike aerosol amount, *aerosol microphysical properties tend to be repeatable* from year to year, for a given source in a given season



Adapted from: Kahn, Survy. Geophys. 2012

Backup Slides

Global Distribution of MISR & MODIS *Coincident*, Retrieved AOD

Overall, **6% to 7%** of overlapping observations produce *coincident*, MISR & MODIS aerosol retrievals


Some coincident coverage over much of the planet

Point density varies

- Desert
- Snow & Ice
- Cloud
- Polar night
- Glint

July 2006

Matched
MISR/MODIS



MISR Only

MODIS Only
(*within* MISR swath)

